



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A1	(11) International Publication Number: WO 97/45555 (43) International Publication Date: 4 December 1997 (04.12.97)
(21) International Application Number: PCT/SE97/00839 (22) International Filing Date: 22 May 1997 (22.05.97) (30) Priority Data: 9602062-3 29 May 1996 (29.05.96) SE (71) Applicant (for all designated States except US): PHARMACIA BIOTECH AB [SE/SE]; S-751 82 Uppsala (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): HASEBE, Masahisa [JP/JP]; 2-2-7, Hattori-dai, Kanmaki-cho, Kitakaturagi-gun, Nora-ken 639-02 (JP). GOTO, Masanori [JP/JP]; 5-11-205, Tamagawa 2-chome, Ota-ku, Tokyo 146 (JP). TOSU, Mariko [JP/JP]; 8-3-1008, Horinouchi 1-chome, Suginami-ku, Tokyo 166 (JP). (74) Agents: EBBINGHAUS, Marie-Louise et al.; Pharmacia & Upjohn AB, Patent Dept., S-751 82 Uppsala (SE).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHOD FOR DETECTION OF MUTATIONS (57) Abstract A method and a kit for detecting a mutation from a non-mutated sequence of a target polynucleotide in a sample, by using a mismatch binding protein. The method comprises: a) providing non-mutated and mutated target polynucleotide; b) forming duplex of the non-mutated and mutated single strands of the target polynucleotide in a); c) adding a single strand binding protein to the polynucleotide from b); d) incubating the mismatch binding protein with an activating agent; e) adding the incubated mismatch binding protein from d) to the polynucleotide from c), whereby the mismatch binding protein binds to the duplex formed by one non-mutated and one mutated single strand of the target polynucleotide; f) detecting the presence of any mismatch binding protein bound to the target polynucleotide. The kit comprises: a) single strand binding protein, b) activator for mismatch binding protein; c) mismatch binding protein; d) amplification primer(s); e) solid support for binding of target polynucleotide or mismatch binding protein.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Method for detection of mutations

The present invention relates to a method and a kit for detecting a mutation from a non-mutated sequence of a target polynucleotide. More specifically the invention relates to a method for detecting mutations by using a mismatch binding protein.

The development of methods for the detection of mutations in DNA is important in diagnostic tests since a large number of human genetic diseases are caused by single base substitutions or by small additions or deletions in the genome. For example, the possibility to detect mutations is an important tool in early detection of cancer.

There are a lot of methods available for detection of mutations. Single strand conformation polymorphism (SSCP) for example, is based on the fact that single stranded DNA that differs by as little as a single base can take up different conformations in a non-denaturing environment, which allows for separation by electrophoresis. In Denaturing gradient gel electrophoresis (DGGE) the altered melting characteristics, exhibited by base pair changes, are used. The melted duplexes are separated when moving into a gel of increasing denaturation concentration. These methods as well as others require gel electrophoresis separation.

Many methods worked out to detect mutations, consisting of one or a few bases, are based on hybridization between a standard DNA (or RNA) and a test DNA. The test sample is mixed with wild-type sample, which can be labelled, and the mixture is denatured and allowed to renature. The mutation is then exposed as a mispaired or unpaired base in a heteroduplex molecule. Different ways are known for the detection of the mispaired or unpaired base. Cleaving of by chemical or enzymatic treatment is one commonly used method. The fragments are observed by gel electrophoresis. Several patent applications describe the use of mismatch binding proteins to detect the heteroduplex molecule (e.g. WO 93/02216, WO 95/12689). DNA mispairing occurs in vivo and is recognized and corrected by a family of repair pro-

teins. Included are proteins which recognize and bind to mismatch containing DNA, so called mismatch binding proteins. It has been found that the mismatch binding proteins forms specific complexes with any of the eight possible mismatched base pairs. The mismatch repair system has been studied in bacteria such as E.coli and Salmonella. MutS is one such protein identified in the E.coli mismatch repair system. The detection of binding of a mismatch binding protein to the DNA is usually made by some type of a label. Recently a rapid method for detection of point mutations using a label free technique was reported (by Ivan Babic et al on the 5th BIA symposium in San Diego, September 1995). The interaction of MutS with short oligonucleotides containing various base-pair mismatches was examined using optical biosensor technology.

The detection methods according to the state of the art based on gel electrophoresis or labelling procedures are time-consuming methods. Therefore, in developing a time-effective assay, these methods are difficult to apply for rapid screening of samples for the detection of mutations. These methods are also difficult to automate.

According to WO 95/12689 and also to Babic et al as mentioned above, MutS binds to DNA containing mispaired bases, but does not bind to DNA without mismatches or single stranded DNA. However, it was found in the present invention that MutS binds non-specifically. Thus, it was found that MutS binds both to single stranded DNA and to double stranded DNA without mismatches. This means that the accuracy of the MutS detection method is not satisfactory for detection of mutations, because non-specific binding of MutS can cause false positive.

The object of the present invention is to obtain an improved method for detecting a mutation by using a mismatch binding protein, without the draw-back mentioned above.

A further object of the invention is to present a kit useful for detection of mutations.

The objects of the invention are achieved by the method and kit as claimed in the claims. According to the inven-

tion a method for detecting a mutation from a non-mutated sequence of a target polynucleotide in a sample is obtained by using a mismatch binding protein. The method comprises:

- a) providing non-mutated and mutated target polynucleotide,
- b) forming duplex of the non-mutated and mutated single strands of the target polynucleotide in a),
- c) adding a single strand binding protein to the polynucleotide from b),
- d) incubating the mismatch binding protein with an activating agent,
- e) adding the incubated mismatch binding protein from d) to the polynucleotide from c), whereby the mismatch binding protein binds to the duplex formed by one non-mutated and one mutated single strand of the target polynucleotide,
- f) detecting the presence of any mismatch binding protein bound to the target polynucleotide.

According to a further aspect of the invention a kit for detecting a mutation from a non-mutated sequence of a target polynucleotide in a sample is achieved. The kit comprises:

- a) single strand binding protein,
- b) activator for mismatch binding protein,
- c) mismatch binding protein,
- d) amplification primer(s),
- e) solid support for binding of target polynucleotide or mismatch binding protein.

With the present invention it was surprisingly found that the non-specific binding of mismatch binding protein could be avoided. Binding of the mismatch binding protein to single strands is inhibited by the single strand binding protein. By activating the mismatch binding protein with an activator, before addition to the sample, binding to double strands lacking mismatches does not take place .

According to a preferred embodiment of the invention, the mutated or the non-mutated target polynucleotide is immobilized on a solid support before the duplex in step b)

is formed. Most preferably the non-mutated target nucleotide is immobilized. Alternatively, the target polynucleotide is immobilized on the solid support after duplex formation in step b) or mismatch binding protein is immobilized on a solid support and binding of the target polynucleotide is detected. According to a further preferred embodiment the solid support is a sensor chip surface. The presence of the mismatch binding protein can then be detected by a label free technique. One such method is detection by surface plasmon resonance (SPR). An optical biosensor system based on SPR is described in WO 90/05295 filed by Pharmacia AB. By this method of detection the time-consuming detection with gel electrophoresis or labelling could be avoided.

In the first step of the invention a sample of the target polynucleotide, i.e. a sequence of a DNA or RNA having a mutation, is mixed with a non-mutated (wild-type) sample of the polynucleotide. The sample mixture is denatured to achieve single strands of the polynucleotide, if the samples are not already single stranded and then the single strands are allowed to renature to double strands. There will then be formed homoduplexes as well as heteroduplexes. A homoduplex is a duplex of two non-mutated strands or two mutated strands. A heteroduplex is a duplex of one non-mutated and one mutated strand. According to one embodiment of the invention the target polynucleotide in the sample can be amplified before the first step of the invention. Conventional amplification methods can be used, especially amplification by PCR (see US 4,683,195; US 4,683,202; US 4,800,159).

In one preferred embodiment a single stranded non-mutated polynucleotide is immobilized to a sensor chip surface and the sample suspecting of containing a mutation of the polynucleotide is added in single stranded form. The added sample is allowed to anneal to the polynucleotide on the sensor chip.

In the next step a single strand binding protein is added which binds to the single strands being left in the

mixture. Before the mismatch binding protein is added to the thus produced sample mixture, the mismatch binding protein is incubated with an activator. It was found that by activating the mismatch binding protein the binding to homoduplexes is prevented. The activated mismatch binding protein is added to the sample mixture and allowed to bind to the heteroduplex. The presence of any bound mismatch protein is then detected.

The immobilization of the polynucleotide is made by conventional methods such as adsorption, covalent binding or some type of a binding pair. One member of that binding pair is attached to the polynucleotide or to the mismatch-binding protein and the other member is applied to the solid support. Example of such binding pairs are biotin - avidin, biotin - streptavidin, cystein - thiol groups, antigen - antibody, lectin - sugar. In a preferred embodiment of the kit the amplification primer(s) or the mismatch binding protein has/have one member of a binding pair attached and the other member of the binding pair is coated on the solid support.

The single strand binding proteins (SSB) used to inhibit the binding of the mismatch protein to single strands could be any of the SSB known in the art. Preferred compounds are single strand binding protein (from *E. coli*, fruit fly, *Xenopus laevis*) and gene 32 protein from T4 bacteriophage and their homologue from other species.

The mismatch binding protein used is one of MutS, HexA, MSH1-6, Rep3, RNaseA, uracil-DNA glycosylase, T4 endonuclease VII, resolvase, a protein which can bind to the mismatch. Before adding the mismatch binding protein to the polynucleotide duplex, the protein is activated with an activating agent. Preferably the activating agent is one of the compounds ATP (adenosine 5'-triphosphate), ADP (adenosine 5'-diphosphate), ATP- γ -S (adenosine 5'-O-(3-thiotriphosphate), AMP-PNP (adenosine 5'-[β , γ -imido]triphosphate) or a nucleotide which can bind to the mismatch binding protein. By this activating of the mismatch binding protein it was found that the non specific

binding to both non-mutated as well as mutated samples was avoided. The activation is made by incubating the mismatch binding protein with the activating agent at ambient temperature for seconds to minutes. Finally the activated protein is added to the polynucleotide mixture.

The detection of the bound mismatch protein can be made by the conventional methods used in the state of the art. However, a preferred method is to use an optical biosensor system according to WO 90/05295, as mentioned above and which is incorporated herein by reference. An optical biosensor provides a simple and easy method to detect mutations and permits automation of systems for clinical applications and diagnostic screening. One such system is BIAcore® (Pharmacia Biosensor AB, Uppsala, Sweden), which is based on WO 90/05295.

The invention will now be illustrated with the following non-limiting examples. With parts and per cent are meant parts by weight and per cent by weight if not differently stated. In the examples the result is presented in figures 1-3 with the meaning:

Figure 1A: A graph of the binding of MutS to oligonucleotides, both double stranded, non-mismatched and mismatched as well as single stranded. The binding is represented in the form of resonance units from the surface plasmon resonance (SPR). MutS is introduced at A and the buffer at B in the figure.

Figure 1B: A graph as in figure 1A but where the binding of MutS to single stranded DNA has been blocked.

Figure 2: A graph of the binding of MutS according to the invention. The binding is represented as in figures 1A and 1B.

Figure 3: The amount of MutS binding for the different kinds of mispairs as shown in table 2 in the sequence listing.

Example 1:

Immobilization and formation of duplex

A sensor chip for optical detection was used. The chip comprises a transparent plate of glass, plastic or another

transparent material. This plate is provided with a metal film on one of its surfaces. A dielectric film is attached to the metal film. On this surface streptavidin is coated.

A biotinylated 20-mer oligonucleotide having the sequence from Val 9 to Gly 15 of N-ras (RB 671A in the sequence listing) at 1 μ M in immobilization buffer comprising 10 mM Tris-HCl, pH 7.5, 0.3 M NaCl, 1mM EDTA, pH 8.0, was flowed over the sensor chip surface at a flow rate of 5 μ l/min at a temperature of 25°C. Then 45 μ l of a μ M 20-mer oligonucleotide having a complementary sequence (681C in the sequence listing) or a single mismatched base (683C in the sequence listing) was flowed over the immobilized oligonucleotide at a flow rate of 5 μ l/min at 25°C to form a double-stranded oligonucleotide on the sensor chip surface by hybridization. The latter double stranded oligonucleotide had G/T mispairing. For the hybridization a buffer containing 0.9 M NaCl and 90 mM sodium citrate was used.

Mismatch detection according to the state of the art

100 nM MutS (E.coli Mismatch Binding Protein from Amersham International) in a binding buffer comprising 50 mM Hepes-KOH, pH 7.2, 100 mM KCl, 1mM EDTA, pH 8.0, 1mM dithiothreitol, 5mM MgCl₂ was flowed over the immobilized complementary or G/T mismatched double stranded oligonucleotide and also over the immobilized single-stranded oligonucleotide at a flow rate of 5 μ l/min at 25°C.

The binding of MutS was detected in a BIAcore® instrument with registration of the SPR. As shown in Figure 1A, the increased curve of the sensorgram expressing the binding of MutS to the DNA indicates an increase in relative resonance units due to binding of MutS not only to the mismatched double stranded DNA but also to the single stranded DNA and the complementary double stranded DNA.

Example 2:

Immobilization and formation of duplex was made in the same manner as in example 1.

Mismatch detection according to the state of the art but modified with single-strand binding protein

45 μ l of 50 μ g/ml of single-strand binding protein
5 (E.coli Single-Strand Binding Protein with mol.weight 19 kDa from Pharmacia Biotech) in binding buffer containing 0.005% Tween20 was flowed over the sensor chip containing immobilized strands at a flow rate of 5 μ l/min at 25°C. MutS was then applied in the same manner as above.

10 The binding of MutS was detected in the same way as in example 1. The resulting sensorgram indicated that single-strand binding protein clearly prevents binding of MutS to single stranded DNA. However, MutS still binds to mismatched and non-mismatched oligonucleotides, see figure 1B.

15 Example 3:

Immobilization and formation of duplex was made in the same manner as in example 1.

Mismatch detection according to the invention

20 The blocking of single stranded DNA was made in the same way as in example 2.

MutS was incubated for 5 minutes at 25°C in the presence of 20mM ATP before use. The ATP incubated MutS was flowed over the immobilized oligonucleotides in the same manner as Fig.1A.

25 The detection in the same way as in example 1 showed that MutS previously incubated with ATP bound only to mismatched DNA and not to non-mismatched DNA, see figure 2.

30 Examples with all the 20 mer oligonucleotides in the sequence list and immobilized and hybridized as in table 2 in the sequence listing, showed that all mismatches could be detected with the method according to the invention, see figure 3.

Sequence Listing

Table 1. Sequences of 20mer oligonucleotides

Name	Sequence
	5' 3'
RB651A Biotin-	ACAGCTGGACAAGAAGAGTA
661T	TACTCTTCTTGTCCAGCTGT
662T	TACTCTTCT <u>A</u> GTCCAGCTGT
664T	TACTCTTCT <u>Q</u> GTCCAGCTGT
665T	TACTCTTCT <u>T</u> GTCCAGCTGT
666T	TACTCTTCTT <u>T</u> GTCCAGCTGT
RB671A Biotin-	GTTGGAGCAGGTGGTGGTGG
681C	CCAACACCACCTGCTCCAAC
682C	CCAACACCAG <u>C</u> TGCTCCAAC
683C	CCAACACCA <u>T</u> CTGCTCCAAC
684C	CCAACACCA <u>A</u> CTGCTCCAAC
689C	CCAACACCT <u>C</u> CTGCTCCAAC

The mismatched bases are underlined in the sequence.

Table 2. Pairs of oligonucleotides to form immobilized dsDNA

dsDNA	Immobilized oligonucleotides	Hybridized oligonucleotides
non-mismatched dsDNA	RB651A	661T
non-mismatched dsDNA	RB671A	681C
A/A mismatched dsDNA	RB651A	662T
A/C mismatched dsDNA	RB651A	664T
C/C mismatched dsDNA	RB651A	665T
T/C mismatched dsDNA	RB651A	666T
G/G mismatched dsDNA	RB671A	682C
G/T mismatched dsDNA	RB671A	683C
G/A mismatched dsDNA	RB671A	684C
T/T mismatched dsDNA	RB671A	689C

Claims

1. A method for detecting a mutation from a non-mutated sequence of a target polynucleotide in a sample, by using a mismatch binding protein, comprising:
- 5 a) providing non-mutated and mutated target polynucleotide,
- b) forming duplex of the non-mutated and mutated single strands of the target polynucleotide in a),
- c) adding a single strand binding protein to the
- 10 polynucleotide from b),
- d) incubating the mismatch binding protein with an activating agent,
- e) adding the incubated mismatch binding protein from d) to the polynucleotide from c), whereby the mismatch
- 15 binding protein binds to the duplex formed by one non-mutated and one mutated single strand of the target polynucleotide,
- f) detecting the presence of any mismatch binding protein bound to the target polynucleotide.
- 20 2. A method according to claim 1, characterized in that before the duplex in step b) is formed, the mutated or the non-mutated target polynucleotide is immobilized to a solid support.
3. A method according to claim 1, characterized in that
- 25 after the duplex is formed in step b) the target polynucleotide is immobilized to a solid support.
4. A method according to claim 1, characterized in that the mismatch binding protein is immobilized on a solid support and the duplex is introduced and binding of the
- 30 target polynucleotide is detected.
5. A method according to claim 2-4, characterized in that the solid support is a sensor chip surface.
6. A method according to claim 5, characterized in that the presence of mismatch binding protein is detected by a
- 35 label free technique.
7. A method according to claim 6, characterized in that the label free technique is surface plasmon resonance.

8. A method according to claim 2, characterized in that the non-mutated target polynucleotide is immobilized on the solid support.

9. A method according to any of the claims 2,3 or 8, characterized in that the polynucleotide is immobilized by a binding pair, one member of the binding pair being attached to the polynucleotide and the other member being coated on the solid support.

10. A method according to claim 4, characterized in that the mismatch binding protein is immobilized by a binding pair, one member of the binding pair being attached to the mismatch binding protein and the other member of the binding pair being coated on the solid support.

11. A method according to claim 1, characterized in that the mismatch binding protein is one of MutS, HexA, MSH1-6, Rep3, RNaseA, uracil-DNA glycosylase, T4 endonuclease VII, resolvase.

12. A method according to claim 1, characterized in that the activating agent in step d) is one of the compounds ATP, ADP, ATP- γ -S, AMP-PNP or a nucleotide which can bind to the mismatch binding protein.

13. A method according to claim 1, characterized in that the target polynucleotide in the sample is amplified before step a).

14. A kit for detecting a mutation from a non-mutated sequence of a target polynucleotide in a sample, said kit comprising

- a) single strand binding protein,
- b) activator for mismatch binding protein,
- c) mismatch binding protein,
- d) amplification primer(s),
- e) solid support for binding of target polynucleotide or mismatch binding protein.

15. A kit according to claim 14, characterized in that the amplification primer(s) or the mismatch binding protein has/have one member of a binding pair attached and the

12

other member of the binding pair is coated on the solid support.

1/4




Fig. 1A

2/4




Fig.1B

3/4




Fig. 2

4/4




Fig. 3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/00839

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C12Q 1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, MEDLINE, BIOSIS, DBA, CA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 9512688 A1 (UNITED STATES BIOCHEMICAL CORPORATION), 11 May 1995 (11.05.95) -----	1-15

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 August 1997

Date of mailing of the international search report

01 -09- 1997

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Patrick Andersson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

06/08/97

International application No.

PCT/SE 97/00839

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9512688 A1	11/05/95	US 5459039 A	17/10/95
		US 5556750 A	17/09/96
		CA 2175490 A	11/05/95
<hr/>			